

# HYPOGLYCAEMIA

**LOW**

Blood Glucose Level < 4.0mmol/L

**DO NOT LEAVE CHILD UNATTENDED  
DO NOT DELAY TREATMENT**

## Signs and symptoms

Note: Symptoms may not always be obvious.  
Pale, headache, shaky, sweaty, dizzy, drowsy, changes in behavior.

### Child conscious

(Able to eat hypo food)

### Step 1: Give fast acting carb

(As supplied or listed on management plan)

### Step 2: Recheck BGL after 15 mins

If BGL < 4.0 repeat Step 1  
If BGL > 4.0 go to Step 3

### Step 3: Give sustaining carb

(As supplied or listed on management plan)

### Child unconscious/drowsy

(Risk of choking/unable to swallow)

### First Aid DRABC

Stay with unconscious child

**CALL AN  
AMBULANCE  
DIAL 000**

### Contact parent/guardian

When safe to do so

PARENT / GUARDIAN NAME \_\_\_\_\_

CONTACT No \_\_\_\_\_

## SCHOOL SETTING

### Multiple daily injections

Use in conjunction with management plan

# DIABETES SCHOOL ACTION PLAN 2017



Photo of child

CHILD'S NAME

\_\_\_\_\_  
\_\_\_\_\_

CENTRE

\_\_\_\_\_  
\_\_\_\_\_

## INSULIN

Insulin is taken 4 or more times per day.

An injection will be needed before lunch.

Able to inject insulin:  independently  
 with supervision  
 with assistance

Injection will be given in: \_\_\_\_\_  
(Room/location)

## ROUTINE BGL CHECKING TIMES

- Anytime, anywhere in the centre
- Prior to lunch
- Anytime hypo is suspected
- Prior to activity
- Prior to exams or tests (e.g. NAPLAN)

## PHYSICAL ACTIVITY

- 1 serve sustaining carb before every 30 mins of activity or swimming
- Vigorous activity should **not** be undertaken if BGL > 15.0 and blood ketones are > 1.0

# HYPERGLYCAEMIA

**HIGH**

Blood Glucose Level > 15.0mmol/L

**HIGH BGLS ARE NOT UNCOMMON**

## Signs and Symptoms

Note: Symptoms may not always be obvious.  
Some could be: increased thirst, increased urine production, poor concentration, irritability, lethargy.

### Child well

Re-check BGL in 2 hours

### Child unwell

E.g. vomiting  
Check blood ketones if able

### Encourage oral fluids, return to class

1-2 glasses water per hour; extra toilet visits may be required

**CONTACT PARENT/  
GUARDIAN  
TO COLLECT  
CHILD ASAP**

**In 2 hours, if BGL still > 15.0  
call parent/guardian for advice**

DATE \_\_\_\_\_

HOSPITAL \_\_\_\_\_

TREATING DNE \_\_\_\_\_

CONTACT No \_\_\_\_\_