## **HYPO**GLYCAEMIA

LOW

Blood Glucose Level < 4.0mmol/L

# DO NOT LEAVE CHILD UNATTENDED DO NOT DELAY TREATMENT

### Signs and symptoms

Note: Symptoms may not always be obvious.

Pale, headache, shaky, sweaty, dizzy, drowsy, changes in behavior.

## Child conscious

(Able to eat hypo food)

# Step 1: Give fast acting carb

(As supplied or listed on management plan)

## Step 2: Recheck BGL after 15 mins

If BGL < 4.0 repeat Step 1 If BGL > 4.0 go to Step 3

# Step 3: Give sustaining carb

(As supplied or listed on management plan)

CALL AN

**AMBULANCE** 

**DIAL 000** 

Child unconscious/

drowsy

(Risk of choking/unable

to swallow)

First Aid DRSABCD

Stav with unconscious

child

## Contact parent/ guardian

When safe to do so

# PARENT / GUARDIAN NAME \_\_\_\_\_\_

### **SCHOOL SETTING**

### **Multiple daily injections**

Use in conjunction with management plan

# **DIABETES SCHOOL ACTION PLAN** 2017

,	CHILD'S NAME
!	
Photo of child	
	CENTRE
1 1 1	

### **INSULIN**

Insulin is taken 4 or more times per day.

An injection will be needed before lunch.

Able to inject insulin: 

independently

with supervision

☐ with assistance
Injection will be given in:

(Room/location)

### **ROUTINE BGL CHECKING TIMES**

- Anytime, anywhere in the centre
- Prior to lunch
- Anytime hypo is suspected
- Prior to activity
- Prior to exams or tests (e.g. NAPLAN)

### PHYSICAL ACTIVITY

- 1 serve sustaining carb before every 30 mins of activity or swimming
- Vigorous activity should **not** be undertaken if BGL >15.0 and blood ketones are >1.0

## **HYPER**GLYCAEMIA

HIGH

Blood Glucose Level > 15.0 mmol/L

### HIGH BGLS ARE NOT UNCOMMON

### **Signs and Symptoms**

Note: Symptoms may not always be obvious. Some could be: increased thirst, increased urine production, poor concentration, irritability, lethargy.

## Child well

Re-check BGL in 2 hours

### Child unwell

E.g. vomiting
Check blood ketones
if able

# Encourage oral fluids, return to class

1-2 glasses water per hour; extra toilet visits may be required CONTACT PARENT/
GUARDIAN
TO COLLECT
CHILD ASAP

In 2 hours, if BGL still >15.0 call parent/guardian for advice

DATE	
HOSPITAL	
TREATING DNE _	
CONTACT No	





